

## POLYNOMIALS

### (A) Main Concepts and Results

Meaning of a Polynomial

Degree of a polynomial

Coefficients

Monomials, Binomials etc.

Constant, Linear, Quadratic Polynomials etc.

Value of a polynomial for a given value of the variable

Zeros of a polynomial

Remainder theorem

Factor theorem

Factorisation of a quadratic polynomial by splitting the middle term

Factorisation of algebraic expressions by using the Factor theorem

Algebraic identities –

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$

$$x^2 - y^2 = (x + y)(x - y)$$

$$(x + a)(x + b) = x^2 + (a + b)x + ab$$

$$(x + y + z)^2 = x^2 + y^2 + z^2 + 2xy + 2yz + 2zx$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3 = x^3 + y^3 + 3xy(x + y)$$

$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3 = x^3 - y^3 - 3xy(x - y)$$

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$$

### (B) Multiple Choice Questions

**Sample Question 1 :** If  $x^2 + kx + 6 = (x + 2)(x + 3)$  for all  $x$ , then the value of  $k$  is

- (A) 1                      (B) -1                      (C) 5                      (D) 3

**Solution :** Answer (C)

### EXERCISE 2.1

Write the correct answer in each of the following :

1. Which one of the following is a polynomial?

(A)  $\frac{x^2}{2} - \frac{2}{x^2}$

(B)  $\sqrt{2x} - 1$

(C)  $x^2 + \frac{3x^2}{\sqrt{x}}$

(D)  $\frac{x-1}{x+1}$

2.  $\sqrt{2}$  is a polynomial of degree

(A) 2

(B) 0

(C) 1

(D)  $\frac{1}{2}$

3. Degree of the polynomial  $4x^4 + 0x^3 + 0x^5 + 5x + 7$  is

(A) 4

(B) 5

(C) 3

(D) 7

4. Degree of the zero polynomial is

(A) 0

(B) 1

(C) Any natural number

(D) Not defined

5. If  $p(x) = x^2 - 2\sqrt{2}x + 1$ , then  $p(2\sqrt{2})$  is equal to

(A) 0

(B) 1

(C)  $4\sqrt{2}$

(D)  $8\sqrt{2} + 1$

6. The value of the polynomial  $5x - 4x^2 + 3$ , when  $x = -1$  is

(A) -6

(B) 6

(C) 2

(D) -2

POLYNOMIALS

15

7. If  $p(x) = x + 3$ , then  $p(x) + p(-x)$  is equal to  
(A) 3 (B)  $2x$  (C) 0 (D) 6
8. Zero of the zero polynomial is  
(A) 0 (B) 1  
(C) Any real number (D) Not defined
9. Zero of the polynomial  $p(x) = 2x + 5$  is  
(A)  $-\frac{2}{5}$  (B)  $-\frac{5}{2}$  (C)  $\frac{2}{5}$  (D)  $\frac{5}{2}$
10. One of the zeroes of the polynomial  $2x^2 + 7x - 4$  is  
(A) 2 (B)  $\frac{1}{2}$  (C)  $-\frac{1}{2}$  (D) -2
11. If  $x^{51} + 51$  is divided by  $x + 1$ , the remainder is  
(A) 0 (B) 1 (C) 49 (D) 50
12. If  $x + 1$  is a factor of the polynomial  $2x^2 + kx$ , then the value of  $k$  is  
(A) -3 (B) 4 (C) 2 (D) -2
13.  $x + 1$  is a factor of the polynomial  
(A)  $x^3 + x^2 - x + 1$  (B)  $x^3 + x^2 + x + 1$   
(C)  $x^4 + x^3 + x^2 + 1$  (D)  $x^4 + 3x^3 + 3x^2 + x + 1$
14. One of the factors of  $(25x^2 - 1) + (1 + 5x)^2$  is  
(A)  $5 + x$  (B)  $5 - x$  (C)  $5x - 1$  (D)  $10x$
15. The value of  $249^2 - 248^2$  is  
(A)  $1^2$  (B) 477 (C) 487 (D) 497
16. The factorisation of  $4x^2 + 8x + 3$  is  
(A)  $(x + 1)(x + 3)$  (B)  $(2x + 1)(2x + 3)$   
(C)  $(2x + 2)(2x + 5)$  (D)  $(2x - 1)(2x - 3)$
17. Which of the following is a factor of  $(x + y)^3 - (x^3 + y^3)$ ?  
(A)  $x^2 + y^2 + 2xy$  (B)  $x^2 + y^2 - xy$  (C)  $xy^2$  (D)  $3xy$
18. The coefficient of  $x$  in the expansion of  $(x + 3)^3$  is  
(A) 1 (B) 9 (C) 18 (D) 27
19. If  $\frac{x}{y} + \frac{y}{x} = -1$  ( $x, y \neq 0$ ), the value of  $x^3 - y^3$  is

- (A) 1                      (B) -1                      (C) 0                      (D)  $\frac{1}{2}$

20. If  $49x^2 - b = \left(7x + \frac{1}{2}\right)\left(7x - \frac{1}{2}\right)$ , then the value of  $b$  is

- (A) 0                      (B)  $\frac{1}{\sqrt{2}}$                       (C)  $\frac{1}{4}$                       (D)  $\frac{1}{2}$

21. If  $a + b + c = 0$ , then  $a^3 + b^3 + c^3$  is equal to

- (A) 0                      (B)  $abc$                       (C)  $3abc$                       (D)  $2abc$

### (C) Short Answer Questions with Reasoning

**Sample Question 1 :** Write whether the following statements are **True** or **False**. Justify your answer.

- (i)  $\frac{1}{\sqrt{5}}x^{\frac{1}{2}} + 1$  is a polynomial                      (ii)  $\frac{6\sqrt{x} + x^{\frac{3}{2}}}{\sqrt{x}}$  is a polynomial,  $x \neq 0$

**Solution :**

- (i) False, because the exponent of the variable is not a whole number.
- (ii) True, because  $\frac{6\sqrt{x} + x^{\frac{3}{2}}}{\sqrt{x}} = 6 + x$ , which is a polynomial.

### EXERCISE 2.2

1. Which of the following expressions are polynomials? Justify your answer:

- (i) 8                      (ii)  $\sqrt{3}x^2 - 2x$                       (iii)  $1 - \sqrt{5}x$
- (iv)  $\frac{1}{5x^{-2}} + 5x + 7$                       (v)  $\frac{(x-2)(x-4)}{x}$                       (vi)  $\frac{1}{x+1}$
- (vii)  $\frac{1}{7}a^3 - \frac{2}{\sqrt{3}}a^2 + 4a - 7$                       (viii)  $\frac{1}{2x}$

2. Write whether the following statements are **True** or **False**. Justify your answer.

- (i) A binomial can have at most two terms
- (ii) Every polynomial is a binomial
- (iii) A binomial may have degree 5
- (iv) Zero of a polynomial is always 0
- (v) A polynomial cannot have more than one zero
- (vi) The degree of the sum of two polynomials each of degree 5 is always 5.

### (D) Short Answer Questions

#### Sample Question 1 :

(i) Check whether  $p(x)$  is a multiple of  $g(x)$  or not, where

$$p(x) = x^3 - x + 1, \quad g(x) = 2 - 3x$$

(ii) Check whether  $g(x)$  is a factor of  $p(x)$  or not, where

$$p(x) = 8x^3 - 6x^2 - 4x + 3, \quad g(x) = \frac{x}{3} - \frac{1}{4}$$

#### Solution :

(i)  $p(x)$  will be a multiple of  $g(x)$  if  $g(x)$  divides  $p(x)$ .

Now,  $g(x) = 2 - 3x = 0$  gives  $x = \frac{2}{3}$

Remainder  $= p\left(\frac{2}{3}\right) = \left(\frac{2}{3}\right)^3 - \left(\frac{2}{3}\right) + 1$

$$= \frac{8}{27} - \frac{2}{3} + 1 = \frac{17}{27}$$

Since remainder  $\neq 0$ , so,  $p(x)$  is not a multiple of  $g(x)$ .

(ii)  $g(x) = \frac{x}{3} - \frac{1}{4} = 0$  gives  $x = \frac{3}{4}$

$g(x)$  will be a factor of  $p(x)$  if  $p\left(\frac{3}{4}\right) = 0$  (Factor theorem)

Now,  $p\left(\frac{3}{4}\right) = 8\left(\frac{3}{4}\right)^3 - 6\left(\frac{3}{4}\right)^2 - 4\left(\frac{3}{4}\right) + 3$

$$= 8 \times \frac{27}{64} - 6 \times \frac{9}{16} - 3 + 3 = 0$$

Since,  $p\left(\frac{3}{4}\right) = 0$ , so,  $g(x)$  is a factor of  $p(x)$ .

**Sample Question 2 :** Find the value of  $a$ , if  $x - a$  is a factor of  $x^3 - ax^2 + 2x + a - 1$ .

**Solution :** Let  $p(x) = x^3 - ax^2 + 2x + a - 1$

Since  $x - a$  is a factor of  $p(x)$ , so  $p(a) = 0$ .

$$\text{i.e., } a^3 - a(a)^2 + 2a + a - 1 = 0$$

$$a^3 - a^3 + 2a + a - 1 = 0$$

$$3a = 1$$

Therefore,  $a = \frac{1}{3}$

**Sample Question 3 :** (i) Without actually calculating the cubes, find the value of  $48^3 - 30^3 - 18^3$ .

(ii) Without finding the cubes, factorise  $(x - y)^3 + (y - z)^3 + (z - x)^3$ .

**Solution :** We know that  $x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)$ .

If  $x + y + z = 0$ , then  $x^3 + y^3 + z^3 - 3xyz = 0$  or  $x^3 + y^3 + z^3 = 3xyz$ .

(i) We have to find the value of  $48^3 - 30^3 - 18^3 = 48^3 + (-30)^3 + (-18)^3$ .

$$\text{Here, } 48 + (-30) + (-18) = 0$$

$$\text{So, } 48^3 + (-30)^3 + (-18)^3 = 3 \times 48 \times (-30) \times (-18) = 77760$$

(ii) Here,  $(x - y) + (y - z) + (z - x) = 0$

$$\text{Therefore, } (x - y)^3 + (y - z)^3 + (z - x)^3 = 3(x - y)(y - z)(z - x).$$

### EXERCISE 2.3

1. Classify the following polynomials as polynomials in one variable, two variables etc.

(i)  $x^2 + x + 1$

(ii)  $y^3 - 5y$

(iii)  $xy + yz + zx$

(iv)  $x^2 - 2xy + y^2 + 1$

2. Determine the degree of each of the following polynomials :

(i)  $2x - 1$

(ii)  $-10$

(iii)  $x^3 - 9x + 3x^5$

(iv)  $y^3 (1 - y^4)$

3. For the polynomial

$$\frac{x^3 + 2x + 1}{5} - \frac{7}{2}x^2 - x^6, \text{ write}$$

(i) the degree of the polynomial

(ii) the coefficient of  $x^3$

(iii) the coefficient of  $x^6$

(iv) the constant term

4. Write the coefficient of  $x^2$  in each of the following :

(i)  $\frac{\pi}{6}x + x^2 - 1$

(ii)  $3x - 5$

(iii)  $(x - 1)(3x - 4)$

(iv)  $(2x - 5)(2x^2 - 3x + 1)$

5. Classify the following as a constant, linear, quadratic and cubic polynomials :

(i)  $2 - x^2 + x^3$

(ii)  $3x^3$

(iii)  $5t - \sqrt{7}$

(iv)  $4 - 5y^2$

(v)  $3$

(vi)  $2 + x$

(vii)  $y^3 - y$

(viii)  $1 + x + x^2$

(ix)  $t^2$

(x)  $\sqrt{2}x - 1$

6. Give an example of a polynomial, which is :

(i) monomial of degree 1

(ii) binomial of degree 20

(iii) trinomial of degree 2

7. Find the value of the polynomial  $3x^3 - 4x^2 + 7x - 5$ , when  $x = 3$  and also when  $x = -3$ .

8. If  $p(x) = x^2 - 4x + 3$ , evaluate :  $p(2) - p(-1) + p\left(\frac{1}{2}\right)$

9. Find  $p(0), p(1), p(-2)$  for the following polynomials :

(i)  $p(x) = 10x - 4x^2 - 3$

(ii)  $p(y) = (y + 2)(y - 2)$

10. Verify whether the following are **True** or **False** :

(i)  $-3$  is a zero of  $x - 3$

(ii)  $-\frac{1}{3}$  is a zero of  $3x + 1$

(iii)  $\frac{-4}{5}$  is a zero of  $4 - 5y$

(iv) 0 and 2 are the zeroes of  $t^2 - 2t$

(v)  $-3$  is a zero of  $y^2 + y - 6$

**11.** Find the zeroes of the polynomial in each of the following :

(i)  $p(x) = x - 4$

(ii)  $g(x) = 3 - 6x$

(iii)  $q(x) = 2x - 7$

(iv)  $h(y) = 2y$

**12.** Find the zeroes of the polynomial :

$$p(x) = (x - 2)^2 - (x + 2)^2$$

**13.** By actual division, find the quotient and the remainder when the first polynomial is divided by the second polynomial :  $x^4 + 1$ ;  $x - 1$

**14.** By Remainder Theorem find the remainder, when  $p(x)$  is divided by  $g(x)$ , where

(i)  $p(x) = x^3 - 2x^2 - 4x - 1$ ,  $g(x) = x + 1$

(ii)  $p(x) = x^3 - 3x^2 + 4x + 50$ ,  $g(x) = x - 3$

(iii)  $p(x) = 4x^3 - 12x^2 + 14x - 3$ ,  $g(x) = 2x - 1$

(iv)  $p(x) = x^3 - 6x^2 + 2x - 4$ ,  $g(x) = 1 - \frac{3}{2}x$

**15.** Check whether  $p(x)$  is a multiple of  $g(x)$  or not :

(i)  $p(x) = x^3 - 5x^2 + 4x - 3$ ,  $g(x) = x - 2$

(ii)  $p(x) = 2x^3 - 11x^2 - 4x + 5$ ,  $g(x) = 2x + 1$

**16.** Show that :

(i)  $x + 3$  is a factor of  $69 + 11x - x^2 + x^3$ .

(ii)  $2x - 3$  is a factor of  $x + 2x^3 - 9x^2 + 12$ .

**17.** Determine which of the following polynomials has  $x - 2$  a factor :

(i)  $3x^2 + 6x - 24$

(ii)  $4x^2 + x - 2$

**18.** Show that  $p - 1$  is a factor of  $p^{10} - 1$  and also of  $p^{11} - 1$ .

**19.** For what value of  $m$  is  $x^3 - 2mx^2 + 16$  divisible by  $x + 2$  ?

**20.** If  $x + 2a$  is a factor of  $x^5 - 4a^2x^3 + 2x + 2a + 3$ , find  $a$ .

**21.** Find the value of  $m$  so that  $2x - 1$  be a factor of  $8x^4 + 4x^3 - 16x^2 + 10x + m$ .



**22.** If  $x + 1$  is a factor of  $ax^3 + x^2 - 2x + 4a - 9$ , find the value of  $a$ .

**23.** Factorise :

(i)  $x^2 + 9x + 18$

(ii)  $6x^2 + 7x - 3$

(iii)  $2x^2 - 7x - 15$

(iv)  $84 - 2r - 2r^2$

**24.** Factorise :

(i)  $2x^3 - 3x^2 - 17x + 30$

(ii)  $x^3 - 6x^2 + 11x - 6$

(iii)  $x^3 + x^2 - 4x - 4$

(iv)  $3x^3 - x^2 - 3x + 1$

**25.** Using suitable identity, evaluate the following:

(i)  $103^3$

(ii)  $101 \times 102$

(iii)  $999^2$

**26.** Factorise the following:

(i)  $4x^2 + 20x + 25$

(ii)  $9y^2 - 66yz + 121z^2$

(iii)  $\left(2x + \frac{1}{3}\right)^2 - \left(x - \frac{1}{2}\right)^2$

**27.** Factorise the following :

(i)  $9x^2 - 12x + 3$

(ii)  $9x^2 - 12x + 4$

**28.** Expand the following :

(i)  $(4a - b + 2c)^2$

(ii)  $(3a - 5b - c)^2$

(iii)  $(-x + 2y - 3z)^2$

**29.** Factorise the following :

(i)  $9x^2 + 4y^2 + 16z^2 + 12xy - 16yz - 24xz$

(ii)  $25x^2 + 16y^2 + 4z^2 - 40xy + 16yz - 20xz$

(iii)  $16x^2 + 4y^2 + 9z^2 - 16xy - 12yz + 24xz$

**30.** If  $a + b + c = 9$  and  $ab + bc + ca = 26$ , find  $a^2 + b^2 + c^2$ .

**31.** Expand the following :

(i)  $(3a - 2b)^3$

(ii)  $\left(\frac{1}{x} + \frac{y}{3}\right)^3$

(iii)  $\left(4 - \frac{1}{3x}\right)^3$

**32.** Factorise the following :

(i)  $1 - 64a^3 - 12a + 48a^2$

$$(ii) \quad 8p^3 + \frac{12}{5}p^2 + \frac{6}{25}p + \frac{1}{125}$$

**33.** Find the following products :

$$(i) \quad \left(\frac{x}{2} + 2y\right)\left(\frac{x^2}{4} - xy + 4y^2\right) \qquad (ii) \quad (x^2 - 1)(x^4 + x^2 + 1)$$

**34.** Factorise :

$$(i) \quad 1 + 64x^3 \qquad (ii) \quad a^3 - 2\sqrt{2}b^3$$

**35.** Find the following product :

$$(2x - y + 3z)(4x^2 + y^2 + 9z^2 + 2xy + 3yz - 6xz)$$

**36.** Factorise :

$$(i) \quad a^3 - 8b^3 - 64c^3 - 24abc \qquad (ii) \quad 2\sqrt{2}a^3 + 8b^3 - 27c^3 + 18\sqrt{2}abc.$$

**37.** Without actually calculating the cubes, find the value of :

$$(i) \quad \left(\frac{1}{2}\right)^3 + \left(\frac{1}{3}\right)^3 - \left(\frac{5}{6}\right)^3 \qquad (ii) \quad (0.2)^3 - (0.3)^3 + (0.1)^3$$

**38.** Without finding the cubes, factorise

$$(x - 2y)^3 + (2y - 3z)^3 + (3z - x)^3$$

**39.** Find the value of

$$(i) \quad x^3 + y^3 - 12xy + 64, \text{ when } x + y = -4$$

$$(ii) \quad x^3 - 8y^3 - 36xy - 216, \text{ when } x = 2y + 6$$

**40.** Give possible expressions for the length and breadth of the rectangle whose area is given by  $4a^2 + 4a - 3$ .

### (E) Long Answer Questions

**Sample Question 1 :** If  $x + y = 12$  and  $xy = 27$ , find the value of  $x^3 + y^3$ .

**Solution :**

$$\begin{aligned} x^3 + y^3 &= (x + y)(x^2 - xy + y^2) \\ &= (x + y)[(x + y)^2 - 3xy] \\ &= 12[12^2 - 3 \times 27] \\ &= 12 \times 63 = 756 \end{aligned}$$

**Alternative Solution :**

$$\begin{aligned}x^3 + y^3 &= (x + y)^3 - 3xy(x + y) \\&= 12^3 - 3 \times 27 \times 12 \\&= 12 [12^2 - 3 \times 27] \\&= 12 \times 63 = 756\end{aligned}$$

**EXERCISE 2.4**

1. If the polynomials  $az^3 + 4z^2 + 3z - 4$  and  $z^3 - 4z + a$  leave the same remainder when divided by  $z - 3$ , find the value of  $a$ .
2. The polynomial  $p(x) = x^4 - 2x^3 + 3x^2 - ax + 3a - 7$  when divided by  $x + 1$  leaves the remainder 19. Find the values of  $a$ . Also find the remainder when  $p(x)$  is divided by  $x + 2$ .
3. If both  $x - 2$  and  $x - \frac{1}{2}$  are factors of  $px^2 + 5x + r$ , show that  $p = r$ .
4. Without actual division, prove that  $2x^4 - 5x^3 + 2x^2 - x + 2$  is divisible by  $x^2 - 3x + 2$ .  
[Hint: Factorise  $x^2 - 3x + 2$ ]
5. Simplify  $(2x - 5y)^3 - (2x + 5y)^3$ .
6. Multiply  $x^2 + 4y^2 + z^2 + 2xy + xz - 2yz$  by  $(-z + x - 2y)$ .
7. If  $a, b, c$  are all non-zero and  $a + b + c = 0$ , prove that  $\frac{a^2}{bc} + \frac{b^2}{ca} + \frac{c^2}{ab} = 3$ .
8. If  $a + b + c = 5$  and  $ab + bc + ca = 10$ , then prove that  $a^3 + b^3 + c^3 - 3abc = -25$ .
9. Prove that  $(a + b + c)^3 - a^3 - b^3 - c^3 = 3(a + b)(b + c)(c + a)$ .